THE HOMOLOGICAL CONTENT OF THE JONES REPRESENTATIONS AT q = -1

JENS KRISTIAN EGSGAARD AND SØREN FUGLEDE JØRGENSEN

ABSTRACT. We generalize a discovery of Kasahara and show that the Jones representations of braid groups, when evaluated at q = -1, are related to the action on homology of a branched double cover of the underlying punctured disk.

As an application, we prove for a large family of pseudo-Anosov mapping classes a conjecture put forward by Andersen, Masbaum, and Ueno [AMU06] by extending their original argument for the sphere with four marked points to our more general case.

Contents

1. Introduction	1
1.1. The Jones representation and homology	2
1.2. Quantum representations	4
1.3. The AMU conjecture	5
2. Jones representations	6
3. Proofs of main theorems	7
3.1. Construction of the (iso)morphism	8
3.2. Proof of equivariance	8
3.3. Injectivity	11
4. The AMU conjecture	12
4.1. Determining stretch factors	15
Appendix A. Examples, experiments, and observations	15
A.1. A pseudo-Anosov in the Torelli group	15
A.2. Infinite order at low levels	16
A.3. Bigelow's element of the kernel of the Burau representation	18
References	18

1. INTRODUCTION

The present paper is concerned with an interpretation of the q-dependent two-row Jones representations of braid groups – discovered in 1983 by Vaughan Jones [Jon83] – at q = -1, in terms of the action of the braid group on the homology of certain double covers of punctured disks and spheres.

Jones' discovery [Jon83] that the representations may be used to define interesting invariants of links, together with Witten's discovery [Wit89] of the relation between these link invariants and the Chern–Simons theory of 3-manifolds, has served as an inspiration for several important mathematical innovations, occasionally collected under the label "quantum topology". Any attempt to include here a full account on these historical developments would necessarily be lacking, and we shall make no such; indeed, for the experts, the Jones representation needs no introduction.

The authors have received funding from the Danish National Research Foundation grant DNRF95 (Centre for Quantum Geometry of Moduli Spaces - QGM). S.F.J. is furthermore supported by the Swedish Research Council Grant 621–2011–3629.

1.1. The Jones representation and homology. Let B_n denote the braid group on n strands, let $\overline{M}(g,n)$ denote the mapping class group of a genus g surface Σ_g^n with n boundary components, and let M(g,m) denote the mapping class group of a genus g surface with m punctures.

Following the notation of Wenzl [Wen88, Thm. 2.2], the Jones representation defines for every Young diagram $\lambda = (\lambda_1 \geq \lambda_2 \geq \cdots \geq \lambda_p)$ with *n* boxes, $\sum_j \lambda_j = n$, a representation π_{λ} of B_n , depending on a parameter *q* which we shall take to be a non-zero complex number which is furthermore assumed to be *n*-regular, i.e. it is not an *l*'th root of unity for any $2 \leq l \leq n$. We shall be dealing primarily with two-row Young diagrams and let $\pi_q^{n,d}$ denote the representation of B_n obtained from $\lambda = (\lambda_1 \geq \lambda_1 - d), d \geq 0$.

Let now *n* be given, and let g = n - 1. There then is a homomorphism $\Psi : B_{2n} \to M(g, 0)$ given by mapping the standard braid generators $\sigma_1, \ldots, \sigma_{2n-1}$ to the (right) Dehn twists about the curves $\gamma_0, \beta_1, \gamma_1, \ldots, \beta_g, \gamma_g$ indicated in Figure 1a respectively; for well-definedness, see e.g. [FM11, Fact. 3.9 and Prop. 3.11]. Similarly, we define $\Psi : B_{2n-1} \to \overline{M}(g, 1)$ by mapping $\sigma_1, \ldots, \sigma_{2n-2}$ to twists about $\gamma_0, \beta_1, \gamma_1, \ldots, \gamma_{g-1}, \beta_g$ respectively (Figure 1b). Notice also that these homomorphisms are exactly those that appear in the Birman–Hilden theorem [BH73] (see also [FM11, Sect. 9.4]).



FIGURE 1. Our naming convention for curves and homology cycles on a surface.

Now, M(g, 0) and $\overline{M}(g, 1)$ act on the corresponding first homology groups by symplectomorphisms with respect to the intersection pairing ω . For m = 0, 1 and $l \ge 1$, we let

$$\tilde{\rho}_{\text{hom}}^{g,l}: \overline{M}(g,m) \to \mathrm{GL}(\Lambda^l H_1(\Sigma_g^m, \mathbb{C})/(\omega \wedge \Lambda^{l-2} H_1(\Sigma_g^m, \mathbb{C})))$$

denote the induced action, i.e. for $\varphi \in \overline{M}(g,m)$ and $v_1, \ldots, v_l \in H_1(\Sigma_a^m, \mathbb{C})$,

$$\tilde{\rho}_{\text{hom}}^{g,l}(\varphi)[v_1 \wedge \dots \wedge v_l] = [(\varphi_*)v_1 \wedge \dots \wedge (\varphi_*)v_l],$$

where we use the conventions that $\Lambda^{-1}H_1(\Sigma_g^m, \mathbb{C}) = \{0\}$ and $\Lambda^0 H_1(\Sigma_g^m, \mathbb{C}) = \mathbb{C}$. Finally, let $\rho_{\text{hom}}^{g,l} = \tilde{\rho}_{\text{hom}}^{g,l} \circ \Psi$ denote the corresponding braid group representations. The study of the Jones representation as contained in this paper was initiated by an attempt

The study of the Jones representation as contained in this paper was initiated by an attempt to generalize the results of [AMU06] to general punctured spheres. Here, the authors show how to relate the representations of the mapping class group of a four times punctured sphere, obtained from the level k quantum representations – closely related to specializations of the parameter q of the Jones representations to certain roots of unity – to an action on the homology of a torus by considering the limit $k \to \infty$.

In [Jon87, Sect. 10], Jones gave explicit matrices for the representation associated to \boxplus (closely related to $\pi_q^{6,0}$, the representation associated to \boxplus). Moreover, the choice of basis is such that all matrix entries are in $\mathbb{Z}[q, q^{-1}]$, so that one obtains this way representations of B_6 for all non-zero values of q rather than only the 6-regular ones. Kasahara [Kas01, Lem. 2.1] observed in the same vein as above that at q = -1, the resulting representation is equivalent to the representation $\rho_{\text{hom}}^{2,2}$. In general, the representation space of $\pi_q^{2n,0}$ has dimension C_{n+1} , the (n + 1)'st Catalan number. Inspired by Kasahara's result and noting the relation

$$C_{n+1} = \binom{2n}{n} - \binom{2n}{n-2},$$

one arrives at the following, which we will prove in Section 3.

Theorem 1.1. The Jones representation $\pi_q^{2n,0}$ has a natural extension to q = -1, for which it is equivalent to $\rho_{\text{hom}}^{g,g}$, where g = n - 1.

As it is well-known that $\pi_q^{2n-1,1}$ is equivalent to $\pi_q^{2n,0}|_{B_{2n-1}}$, the above Theorem allows us to deduce a similar homological description for $\pi_q^{2n-1,1}$. On the other hand, $\pi_q^{2n-1,2n-3}$ is nothing but the reduced Burau representation, which at

On the other hand, $\pi_q^{2n-1,2n-3}$ is nothing but the reduced Burau representation, which at q = -1 is known to be equivalent to the representation $\rho_{\text{hom}}^{g,1}$ on $H_1(\Sigma_g^1, \mathbb{C})$, g = n - 1. This suggests that the intermediary representations $\pi_q^{2n-1,d}$ interpolate at q = -1 between appropriate exterior powers of homological representations, and the following result shows that this is indeed the case.

Theorem 1.2. The representation $\pi_q^{2n+1,d}$ has a natural extension to q = -1 for which it is equivalent to the action $\rho_{\text{hom}}^{g,l}$, where g = n and 2l = 2n + 1 - d.

This leaves us with the two-row diagrams having an even number of boxes. The fact that $\pi_q^{n,d}|_{B_{n-1}} \cong \pi_q^{n-1,d-1} \oplus \pi_q^{n-1,d+1}$ (see e.g. [Jon87]), together with the observation that

$$\binom{2g}{l} - \binom{2g}{l-2} + \binom{2g}{l+1} - \binom{2g}{l-1} = \binom{2g+1}{l+1} - \binom{2g+1}{l-1}$$

for all g and l, leads us to the following: let $\hat{\rho}_{\text{hom}}^{g,l}$ denote the action of B_{2n} on $\Lambda^l H_1(\Sigma_g^2, \mathbb{C})$, where g = n - 1, given by mapping $\sigma_1, \ldots, \sigma_{2n-1}$ to the action induced by the homological action of the Dehn twists $t_{\gamma_0}, t_{\beta_1}, t_{\gamma_1}, \ldots, t_{\beta_g}, t_{\gamma_g}$ respectively (see Figure 1c).

Theorem 1.3. The representation $\pi_q^{2n,d}$ has a natural extension to q = -1 for which it is equivalent to a subrepresentation of $\hat{\rho}_{hom}^{g,l}$, where g = n - 1 and 2l = 2n - d.

In each of the three cases, the appropriate intertwining operator is constructed as follows: a natural basis for the representation spaces of $\pi_a^{n,d}$ is given in terms of non-intersecting paths in the

relevant punctured disc, connecting the punctures. Regarding the punctures as marked points, the disk is realized as the quotient of a surface by the order two element rotating the surface by π along its horizontal axis in Figure 1, allowing us to realize the surface as a well-understood branched double cover. Lifting the non-intersecting paths through the double cover defines a collection of loops in the covering surface, and by taking an appropriately ordered and scaled wedge product of the homology classes of these loops, we obtain our desired linear map.

As the insightful reader has no doubt pondered, the exact choice of curves in the double cover whose twists we have chosen to lift the B_n -action to is of little importance, and for Theorems 1.2 and 1.3 any linearly independent set of appropriately intersecting curves would do the job. A similar statement is true for Theorem 1.1; the exact conditions will be made precise in Section 3.

1.2. Quantum representations. Let $\Lambda_{N,k}$ be the set of Young diagrams of height strictly less than N and length at most k. This set comes with an involution [†], which assigns to a Young diagram ($\lambda_1 \geq \cdots \geq \lambda_p$) the result of taking the complement of λ in the Young diagram consisting of N rows, each containing λ_1 boxes, and rotating the result 180 degrees (see [Bla00, p. 206]).

Let Σ be a surface and P a finite set of pairs (p, v) where $p \in \Sigma$ and $v \in (T_p \Sigma \setminus \{0\})/\mathbb{R}_+$, and let $\lambda : P \to \Lambda_{N,k}$ be a labelling of P. Let $\widetilde{M}(\Sigma, P, \lambda)$ be the set of isotopy classes of diffeomorphisms of Σ preserving P and respecting the labelling. The *level k quantum* SU(N)-representation assigns to (Σ, P, λ) a finite dimensional vector space $V_{N,k}(\Sigma, P, \lambda)$ and a projective representation $\rho_{N,k}$ of $\widetilde{M}(\Sigma, P, \lambda)$ on $V_{N,k}$.

These quantum representations, arising originally from the mathematical realizations of Witten's 2+1 dimensional topological quantum field theory constructed from Chern–Simons theory [Wit89], have been studied from a number of different perspectives, including that of the representation theory of quantum groups [RT90], [RT91], [Tur10], the skein theory of the Jones polynomial [BHMV92], [BHMV95], [Bla00], conformal field theory [TUY89], [Uen08], [Kan89], [AU07b], and the geometric quantization of moduli spaces of flat connections [Hit90], [And06]. For a complete description of the equivalence between these constructions, we refer to [AU07a], [AU07b], [AU12] and [AU11]. While useful to have in mind, the present work is independent of this equivalence.

Indeed, for our purposes, it will be convenient to take as our definition of quantum representations those that are obtained from the general procedure, described in [Tur10, Sect. IV.5], which defines quantum representations from modular functors, which in turn we apply to the modular functor arising from conformal field theory, constructed in [AU07b]. The label set of this theory is a certain k-dependent set of dominant integral $\mathfrak{sl}(N, \mathbb{C})$ -weights which is in natural correspondence with the set $\Lambda_{N,k}$ defined above (see [AU11, Lemma 7.1]).

In [Tur10, Sects. IV.4–9], the projective ambiguity of the quantum representations is discussed in detail. We record here the fact that if Σ has genus 0, the quantum representations define not just projective representations, but actual representations. To be concrete, we consider $\Sigma = S^2 = \mathbb{C} \cup \{\infty\}$, and let the set P consist of the points $\{1, \ldots, n, \infty\}$, all framed along the real axis. Label the n + 1 points by $\Box, \ldots, \Box, \lambda^{\dagger}$, where λ belongs to the set $L_{N,k} \subseteq \Lambda_{N,k}$ defined as follows: as in [Wen88, Def. 2.4], we denote by $\Lambda_n^{(N,k+N)}$ the set of Young diagrams with n boxes, at most Nrows, and satisfying furthermore that $\lambda_1 - \lambda_N \leq k$. Let $T : \Lambda_n^{(N,k+N)} \to \Lambda_{N,k}$ be the map which removes all columns of height N, and let

(1)
$$L_{N,k} = T(\{\lambda \in \Lambda_n^{(N,k+N)} \mid (\lambda_2 > 0, \, \lambda_3 = 0) \text{ or } (\lambda_1 > 1, \, \lambda_2 = 1)\}).$$

That is, for k large enough, $L_{N,k}$ consists of those diagrams obtained by applying T to 2-row diagrams and hooks respectively; out of these, the former collection of diagrams will be the most interesting to us.

We write $\widetilde{M}(0,n)_{\infty}$ for the subgroup of $\widetilde{M}(\Sigma, P, (\Box, \ldots, \Box, \lambda^{\dagger}))$ consisting of classes of diffeomorphisms fixing ∞ and its framing. Moreover, we introduce $\widetilde{M}(0,n) = \widetilde{M}(\Sigma, \tilde{P}, (\Box, \ldots, \Box))$, where $\tilde{P} = P \setminus \{\infty\}$.

Now, $\widetilde{M}(0,n)_{\infty}$ is the ribbon braid group on *n* ribbons, which contains the ordinary braid group B_n as a subgroup, by assigning the blackboard framing to a braid. We will denote by $\rho_{N,k}^{\lambda}$ the restriction of the quantum representation of $\widetilde{M}(0,n)_{\infty}$ to this B_n , and will think of B_n as

 $M(0,n)_{\infty}$, the mapping class group of a sphere with n marked points and an extra marked point called ∞ which also carries a preserved projective tangent vector. Finally, we denote by $\rho_{N,k}$ the quantum representation of $\widetilde{M}(0,n)$.

1.3. The AMU conjecture. The AMU conjecture [AMU06, Conjecture 2.4] is concerned with the images of pseudo-Anosovs in the quantum SU(N)-representations of mapping class groups for large values of k. As mentioned above, in [AMU06] the authors prove their conjecture for pseudo-Anosovs in M(0, 4) by relating a limit of quantum representation to homology, and it is this argument that we seek to generalize in the present paper. With Theorem 1.4 below, we provide a large family of mapping classes for which the conjecture holds; previously, the conjecture has only been studied for M(0, 4) in [AMU06], and for M(1, 1) in the closely related quantum SO(3)representations in unpublished work by Masbaum [AMU06, Rem. 5.9], [Mas08], and in [San12].

More precisely, we apply Theorem 1.1 to prove the AMU conjecture for those pseudo-Anosov mapping classes whose dynamics are described in homological terms; one particular family for which this is the case are the ones we will refer to as homological pseudo-Anosovs. Here, we say that $\varphi \in M(0, n)_{\infty}$ is a homological pseudo-Anosov if its image in M(0, n + 1) – under the map forgetting the tangent vector at ∞ – admits invariant transverse measured singular foliations having only even-pronged non-puncture singularities and only odd-pronged puncture singularities away from ∞ . Note that we will systematically make the common abuse of notation and think of φ as both a mapping class and a pseudo-Anosov representing it.

Theorem 1.4. Let $\varphi \in M(0, n)_{\infty}$ be a homological pseudo-Anosov, and let $\lambda \in L_{N,k}$ as given in (1). Then $\rho_{N,k}^{\lambda}(\varphi)$ has infinite order for all but finitely many k.

Remark 1.5. Note that the assumption that $\lambda \in L_{N,k}$ may likely be dropped as long as the trivial cases are omitted. Moreover, as previously mentioned, the 2-diagram case is really the one of interest to us, and the hook case will follow by similar – but easier – reasoning. Note also that there is a well-known duality between Jones representations associated to transpose Young diagrams (see [Jon87, Rem. 3.4] and more generally [FLW02, Thm. 3.1]); taking this into account we could enlarge slightly the set $L_{N,k}$.

As above, we say that $\varphi \in M(0, n)$ is a *homological pseudo-Anosov* if its image in M(0, n) is a pseudo-Anosov whose invariant foliations have the property that all non-puncture singularities are even-pronged and all punctures have odd-pronged singularities.

Theorem 1.6. Assume that n is even. If $\varphi \in \widetilde{M}(0,n)$ is a homological pseudo-Anosov, then $\rho_{2,k}(\varphi)$ has infinite order for all but finitely many k.

Finally, let us mention one way our main theorem may be used to deal with general, not necessarily homological, pseudo-Anosovs.

Corollary 1.7. Let f be a representative for $[f] \in M(0,n)_{\infty}$ whose image in M(0,n+1) is pseudo-Anosov, and let P denote the set of singularities of the (un)stable foliation for f with an odd number of prongs. Then f defines an element in $\widetilde{M}(S^2, |P|)_{\infty}$, and the image of [f] under the quantum SU(N)-representation – where each point of P is labelled by \Box , and ∞ is labelled by any $\lambda \in L_{N,k}$ – has infinite order for all but finitely many values of k.

Likewise, if f represents a pseudo-Anosov $[f] \in M(0,n)$ with P the set of odd-pronged singularities, then f defines an element of M(0, |P|) whose image under $\rho_{2,k}$ has infinite order for all but finitely many levels.

Strictly speaking, in order to deal with the issue of framings, we defined only the genus 0 quantum representations in the case where the marked points were $1, \ldots, n$ and possibly ∞ , so to be precise we should include in the statement a conjugation by a group isomorphism identifying the mapping class groups; this does not change the conclusion.

Proof of Corollary 1.7. The first claim follows directly from Theorem 1.4. By the Euler–Poincaré formula, a singular foliation must have an even number of singularities with an odd number of prongs, and so the second claim follows from Theorem 1.6. \Box



FIGURE 2. The multiplication in TL_4



FIGURE 3. The element $e_3 \in TL_6$

Moreover, we show in Corollary 4.9 that for the homological pseudo-Anosovs, the quantum representations determine their stretch factors, answering positively [AMU06, Question 1.1 (2)] in this case. In Appendix A we discuss the extent to which this is true for non-homological pseudo-Anosovs. As a different spin-off, we show in Proposition A.2 that there are quantum SU(2)-representations whose image at level 8 is infinite; this extends an earlier result due to Masbaum, [Mas99].

Finally, we should note that the related problem of giving homological interpretations of the Burau representation of braids for various values of q in terms of finite covers, and in particular its relation to determining stretch factors of pseudo-Anosovs via representations, has been studied by a number of people; see e.g. [BB07], [McM], [Kob12] and the references contained in these papers.

Acknowledgements. We would like to thank Jørgen Ellegaard Andersen for first introducing us to the conjectures of [AMU06], as the attempt to generalize the results of [AMU06] is what led us to the discoveries in this paper.

2. Jones representations

In this section, we briefly recall the definition of the Jones representations of braid groups, fixing the notation and normalizations used throughout the rest of the paper.

The Temperley-Lieb algebra TL_n is an algebra over $\mathbb{C}(A)$ and has a basis consisting of noncrossing pairings of 2n points, n of them located at the bottom of a square and n of them at the top. The multiplication is given by stacking two squares on top of each other, rescaling the vertical direction to obtain a square, removing all circles, and multiplying by $-A^2 - A^{-2}$ for each removed circle, see Figure 2. It is generated, as an algebra, by n elements, $\operatorname{id}, e_1, e_2, \ldots, e_{n-1}$ (Figure 3). Following Jones, we define a representation of the braid group B_n by $\sigma_i \mapsto \operatorname{Aid} + A^{-1}e_i$. There exists a Markov trace on TL_n and the trace of the Jones representation is essentially the Kauffman bracket of the closure of the braid.

For generic A, the irreducible sectors of this braid group action are in correspondence with $d \in \{0, \ldots, n\}$ with $d \equiv n \mod 2$, and each of the underlying vector spaces has a basis given by trivalent trees with n + 1 leaves, coloured admissibly by Jones–Wenzl idempotents, such that n leaves are coloured by 1 and the remaining leaf by d. This description is well-known to experts, but for concrete and concise reference, see e.g. [Wan10, Thm. 1.20, Thm. 1.23]; indeed the treatment contained in [Wan10, Sect. 1] completely covers what we need.

It is possible to specialize the value of A to a non-zero complex number, obtaining the \mathbb{C} -algebra $\operatorname{TL}_n(A)$, and as long as A is not a 4k+8'th root of unity for any k < n, this algebra is semi-simple, and the above discussion provides us with representations $\rho_A^{n,d} : B_n \to W_A^{n,d}$ (we remark that in the notation of [Wan10, Sect. 1.1.6], $W_A^{n,d} = \operatorname{Hom}(d, 1^n)$, and the basis elements described above are denoted $\{e_{C,d}\}$).

Lemma 2.1. Let $A \in \mathbb{C}$ be as above, and let $q = A^4$. Then the braid group representation $\sigma_i \mapsto A^{-1}\rho_A^{n,d}(\sigma_i)$ is equivalent to $\pi_q^{n,d}$.



FIGURE 4. A basis element of $V^{6,4}$.



FIGURE 5. The actions of σ_2 , σ_3 , and σ_4 on the basis element from Figure 4.

Proof. Once again, this is well-known to experts. A basis of the representation space for $\pi_q^{n,d}$ is given by standard Young tableaux on the relevant Young diagram, to each such tableau there is naturally associated a coloured tree in $W_A^{n,d}$, and this association is bijective. Thus, showing the equivalence boils down to checking that the actions on each of the bases agree under this correspondence. For $\pi_q^{n,d}$, this action is given explicitly in [Wen88, (2.3)], and for $\rho_A^{n,d}$ as explicitly on [Wan10, p. 18].

Recall that our goal is to manipulate the $\pi_q^{n,d}$ into representations that we may specialize to all non-zero values of q, rather than just the *n*-regular ones. With an eye towards Wenzl's recursive formula for the Jones–Wenzl idempotents, one finds that all problematic factors occur in the terms that are not of "highest order" (in the language of [FWW02]). This leads us to the following:

Definition 2.2. Let $V^{n,d}$, $0 \le d \le n$, denote the complex vector space spanned by non-crossing pairings of n + d points, such that n points are "at the top" and d points "at the bottom", in such a way that a point at the bottom will be connected to a point at the top; see Figure 4.

The braid group B_n acts on this space exactly as on the Temperley–Lieb algebra: by stacking braids on diagrams, and resolving the result by using the Kauffman bracket. However, if two bottom points end up connected when doing so, the result is multiplied by 0; see Figure 5.

Lemma 2.3. The representation $\eta_A^{n,d}$ of B_n on $V^{n,d}$ is equivalent to $\rho_A^{n,d}$ when $A \in \mathbb{C} \setminus \{0\}$ is not a 4k + 8 th root of unity for any k < n.

Proof. Recall that $W_A^{n,d}$ has a basis given in terms of trees coloured by Jones–Wenzl idempotents. Replacing an idempotent by the corresponding multicurve, ignoring lower order terms, we obtain a map $W_A^{n,d} \to V^{n,d}$, which is B_n -equivariant by construction. By assumption on A, the k-admissibility condition of coloured trees (c.f. [Wan10, Def. 1.24]) boils down to the triangle inequality, which in turn means that the map is surjective.

Since $\eta_A^{n,d}$ is defined over $\mathbb{Z}[A, A^{-1}]$, we obtain our desired extension of $\pi_q^{n,d}$ by letting $q = A^4$ and taking the tensor product with the 1-dimensional representation $\sigma_i \mapsto A$.

We should note that considering the representation in this particular basis is not a new idea: see for instance [RSA12] for a thorough description of the action which moreover puts focus on the case q = -1 which is of interest to us.

3. Proofs of main theorems

In this section, we show Theorems 1.1, 1.2, and 1.3. Throughout the section, we assume that $A = \exp(-\pi i/4)$.



FIGURE 6. The arcs contributing to w(e) = 3 have been coloured red, and the arcs contributing to v(e) = 2 have been coloured blue.

3.1. Construction of the (iso)morphism. We now construct the intertwining morphism φ as needed in Theorems 1.2 and 1.3. For Theorem 1.1 the construction needs to be tweaked slightly; a concern we defer until it becomes relevant.

For a diagram D in the basis of $V^{n,d}$ we denote by D_0 the set of arcs connecting two top points. We identify the points at the top with the numbers 1, 2, ..., n, and for $a \in D_0$ we denote by a_0 and a_1 the left and right end point of a respectively. The set of bottom points will be called ∞ . The symbol *i* will be used to denote indices and the imaginary unit interchangeably; this should cause no confusion.

Definition 3.1. Let c_1, \ldots, c_{n-1} be simple closed curves on the surface Σ_g^m , such that $c_i \cap c_{i+1}$ consists of a single point, and such that c_i and c_j are disjoint for |i-j| > 1. We orient the curves such that $t_{c_i}(c_{i+1}) = c_{i+1} + c_i$, where we abuse notation and use c_i also for the element that the oriented curve defines in $H_1(\Sigma_q^m, \mathbb{C})$. Assume that span $\{c_i \mid i = 1, \ldots, n-1\}$ has dimension n-1.

We order the arcs in D_0 by their starting points so that e < e' if $e_0 < e'_0$. To each arc e in D_0 we associate the element $X_e = \sum_{i=e_0}^{e_1-1} c_i \in \operatorname{span}\{c_i\} \subseteq H_1(\Sigma_g^m, \mathbb{C}).$ Define a map $\varphi: V^{n,d} \to \Lambda^l H_1(\Sigma_g^m, \mathbb{C})$, by letting, for a basis diagram $D \in V^{n,d}$,

(2)
$$\varphi(D) = f(D) \bigwedge_{e \in D_0} X_e,$$

where $f(D) = (-i)^{\sum_{e \in D_0} w(e) + v(e)}$; here, w(e) denotes the number of arcs between the starting and ending point of e, v(e) denotes the number of points greater than e_1 that are connected to ∞ (see Figure 6), and the wedge product runs through D_0 from the first to the final arc with respect to the ordering on D_0 .

3.2. Proof of equivariance.

Lemma 3.2. Denote by T_i the action of e_i on $V^{n,d}$. Then we have, for all i = 1, 2, ..., n-1 and $D \in V^{n,d}$, that

$$\varphi(D + iT_iD) = t_{c_i}\varphi(D).$$

Proof. Let $c = c_i$. We first observe that $(t_c)_*$ acts trivally on X_e if e is not connected to exactly one of i or i + 1. In particular, if the points i, i + 1 are connected, or if both are connected to ∞ , then

$$(t_c)_*(\varphi(D)) = \varphi(D).$$

Since in these cases $T_i D = 0$, we obtain the claim of the Lemma.

Let us therefore consider the case where we have two distinct arcs connecting to i, i + 1, not both connecting to ∞ . Let us first assume that i+1 is connected to ∞ , and let us denote the arc ending in i by a.

Then $(t_c)_*$ acts trivially on all factors of $\varphi(D)$ except for X_a . If a is the j'th arc in D_0 , then

$$(t_c)_*\varphi(D) = f(D)X_1^{j-1} \wedge (X_a - c) \wedge X_{j+1}^k = \varphi(D) + f(D)X_1^{j-1} \wedge (-c) \wedge X_{j+1}^l$$

where we denote the *i*'th edge in D_0 by e^i and write

$$X_i^j = \bigwedge_{l=i}^j X_{e^i}.$$

On the other hand, we have

$$\begin{split} \varphi(T_i(D)) &= (-i)^{-2w(a)-1} f(D) X_1^{j-1} \wedge X_{j+1}^{j+w(a)} \wedge c \wedge X_{j+w(a)+1}^l \\ &= (-i)^{-2w(a)-1} (-1)^{w(a)} f(D) X_1^{j-1} \wedge c \wedge X_{j+1}^l \\ &= -i f(D) X_1^{j-1} \wedge (-c) \wedge X_{j+1}^l. \end{split}$$

Likewise, if i is connected to ∞ but i + 1 is not, we denote by a the arc connecting to i + 1 and find that

$$(t_c)_*\varphi(D) = f(D)X_1^{j-1} \wedge (X_a + c) \wedge X_{j+1}^l = \varphi(D) + f(D)X_1^{j-1} \wedge c \wedge X_{j+1}^l,$$

and

$$\varphi(T_i(D)) = (-i)^{w(a)-w(a)+1} f(D) X_1^{j-1} \wedge c \wedge X_{j+1}^l = -if(D) X_1^{j-1} \wedge c \wedge X_{j+1}^l.$$

Now only the cases where two distinct arcs from D_0 connect to i and i + 1 remain. Denote these arcs by a and b, and assume that they are the j'th and m'th arc respectively, with j < m. There are three possibilities, corresponding to whether or not a and b start or end in i and i+1 (the case where a starts in i and b ends in i+1 is not possible if $a \neq b$). We denote these cases by the signs (ξ_a^i, ξ_b^i) of the intersections of the curves with c, i.e. $(t_c)_* X_a = X_a + \xi_a^i c$, and likewise for b. Then

$$\begin{aligned} (t_c)_*\varphi(D) &= f(D)X_1^{j-1} \wedge (X_a + \xi_a^i c) \wedge X_{j+1}^{m-1} \wedge (X_b + \xi_b^i c) \wedge X_{m+1}^l \\ &= \varphi(D) + f(D)X_1^{j-1} \wedge (\xi_b^i X_a - \xi_a^i X_b) \wedge X_{j+1}^{m-1} \wedge c \wedge X_{m+1}^l \\ &= \varphi(D) + f(D)X_1^{j-1} \wedge (\xi_b^i X_a \pm c - \xi_a^i X_b) \wedge X_{j+1}^{m-1} \wedge c \wedge X_{m+1}^l \end{aligned}$$

If now $(\xi_a, \xi_b) = (-, +)$, then we have

$$\varphi(T_i(D)) = (-i)^{-w(a)-w(b)+w(a)+w(b)+1} f(D) X_1^{j-1} \wedge (a+c+b) \wedge X_{j+1}^{m-1} \wedge c \wedge X_{m+1}^l$$
$$= -if(D) X_1^{j-1} \wedge (a+c+b) \wedge X_{j+1}^{m-1} \wedge c \wedge X_{m+1}^l.$$

If $(\xi_a, \xi_b) = (+, +),$

$$\begin{split} \varphi(T_i(D)) &= (-i)^{-w(a)-w(b)+w(a)-w(b)-1} f(D) \\ &\quad \cdot X_1^{j-1} \wedge c \wedge X_{j+1}^{m-1} \wedge X_{m+1}^{m+w(b)} \wedge (a-c-b) \wedge X_{m+w(b)+1}^l \\ &= i(-i)^{-2w(b)} (-1)^{w(b)} f(D) X_1^{j-1} \wedge c \wedge X_{j+1}^{m-1} \wedge (a-c-b) \wedge X_{m+1}^l \\ &= -if(D) X_1^{j-1} \wedge (a-c-b) \wedge X_{j+1}^{m-1} \wedge c \wedge X_{m+1}^l. \end{split}$$

And finally, if $(\xi_a, \xi_b) = (-, -),$

$$\begin{aligned} \varphi(T_i(D)) &= (-i)^{-w(a)-w(b)+w(a)-w(b)-1} f(D) \\ &\cdot X_1^{j-1} \wedge (a-c-b) \wedge X_{j+1}^{m-1} \wedge X_{m+1}^{m+w(b)} \wedge c \wedge X_{m+w(b)+1}^l \\ &= i(-i)^{-2w(b)} (-1)^{w(b)} f(D) X_1^{j-1} \wedge c \wedge X_{j+1}^{m-1} \wedge (a-c-b) \wedge X_{m+1}^l \\ &= -if(D) X_1^{j-1} \wedge (-a+c+b) \wedge X_{j+1}^{m-1} \wedge c \wedge X_{m+1}^l. \end{aligned}$$

This shows that φ is a homomorphism between the representation $\sigma_i \mapsto A^{-1}\eta_A^{n,d}(\sigma_i)$ and the representation on $\Lambda^l H_1(\Sigma_g^m, \mathbb{C})$ in the cases of Theorems 1.2, and 1.3. For Theorem 1.1, we define $\tilde{\varphi}(D)$ for $D \in V^{2n,0}$ in the following way: there is an *i* such that there is an arc *a* connecting *i* to *n*. We define an element \tilde{D} in $V^{n-1,1}$ by removing *a*, forgetting *n*, and connecting *i* to ∞ , and we define $\tilde{\varphi}: V^{2n,0} \to \Lambda^g H_1(\Sigma_g, \mathbb{C})$ by $\tilde{\varphi}(D) = \varphi(\tilde{D})$, identifying $H_1(\Sigma_g, \mathbb{C})$ with $H_1(\Sigma_g^1, \mathbb{C})$. It is clear that the induced map $\tilde{\cdot}: V^{2n,0} \to V^{2n-1,1}$ is an isomorphism of vector spaces.

Lemma 3.3. The map $\tilde{\varphi}$ is a homomorphism of representations.

In order to prove this, we need the following Lemma.



FIGURE 7. Here, the outermost arc is a and the b_k 's have been coloured red.

Lemma 3.4. If D is a diagram and a is an arc that connects i to j, i < j, we have that

$$X_a \wedge \bigwedge_{e \prec a} X_e = \left(\sum_{\substack{i \le k \le j, \\ k \equiv i \mod 2}} c_k\right) \wedge \bigwedge_{e \prec a} X_e,$$

where we say that $e \prec a$ if $a_0 < e_0 < e_1 < a_1$.

Proof. The proof is by induction on w(a). If w(a) = 0, the claim is clear. Otherwise we use the induction hypothesis on the arcs b_k , k = 1, 2, ..., m, connecting (i_k, j_k) , where $i_1 = i + 1$, $j_m = j - 1$, and $j_k + 1 = i_{k+1}$ for k < m. See Figure 7.

Up until this point, we have not used the possible description of the curves c_i in terms of the curves β_i and γ_i of Figure 1, but for the proof of Lemma 3.3 below, we will need that

$$c_{n-1} = -\sum_{\substack{1 \le k \le n-3, \\ k \equiv 1 \mod 2}} c_k$$

in $H_1(\Sigma_q, \mathbb{C})$, which will be guaranteed by considering the concrete curves.

Proof of Lemma 3.3. If we consider only the action of $B_{n-1} \subseteq B_n$, then $\tilde{\varphi}$ defines a homomorphism by the above, so we only have to check the equivariance of the action of σ_{n-1} . If n-1 and n are connected, then β_g does not appear in $\varphi(\tilde{D})$, and the action of σ_{n-1} on homology is trivial, just as $T_n(D) = 0$.

If n is connected to i < n - 1 by the j'th arc, and n - 1 is an end point of the m'th arc, which we denote by b, then

$$\begin{split} (t_{c_{n-1}})_*(\varphi(\tilde{D})) &= f(\tilde{D})X_1^{j-1} \wedge X_{j+1}^{m-1} \wedge (X_b - c_{n-1}) \wedge X_{m+1}^l \\ &= \varphi(\tilde{D}) - f(\tilde{D})X_1^{j-1} \wedge X_{j+1}^{m-1} \wedge c_{n-1} \wedge X_{m+1}^l \\ &= \varphi(\tilde{D}) + f(\tilde{D})X_1^{j-1} \wedge X_{j+1}^{m-1} \wedge \left(\sum_{\substack{1 \le k \le n-3, \\ k \equiv 1 \mod 2}} c_k\right) \wedge X_{m+1}^l \\ &= \varphi(\tilde{D}) + f(\tilde{D})X_1^{j-1} \wedge X_{j+1}^{m-1} \wedge \left(\sum_{\substack{i \le k \le b_0, \\ k \equiv 1 \mod 2}} c_k\right) \wedge X_{m+1}^l \end{split}$$

Here we used Lemma 3.4 in last equality, applying it to the arcs left of i that are not contained between the end points of any other arc, and on the arcs between the end points of b. Letting pbe the number of arcs to the right of i in \tilde{D} , we find, where we denote by d an arc connecting i with b_0 , that

$$\begin{split} \widetilde{\varphi(T_{n-1}(D))} &= (-i)^{p-w(b)+p-w(b)-1} f(\tilde{D}) X_1^{j-1} \wedge X_d \wedge X_{j+1}^{m-1} \wedge X_{m+1}^l \\ &= (-i)^{p-w(b)+p-w(b)-1} f(\tilde{D}) X_1^{j-1} \wedge \left(\sum_{\substack{i \le k \le b_0, \\ k \equiv 1 \mod 2}} c_k\right) \wedge X_{j+1}^{m-1} \wedge X_{m+1}^l \\ &= (-1)^{p-w(b)+1} (-i)^{p-w(b)+p-w(b)-1} f(\tilde{D}) X_1^{j-1} \wedge X_{j+1}^{m-1} \wedge \left(\sum_{\substack{i \le k \le b_0, \\ k \equiv 1 \mod 2}} c_k\right) \wedge X_{m+1}^l \\ &= -if(\tilde{D}) X_1^{j-1} \wedge X_{j+1}^{m-1} \wedge \left(\sum_{\substack{i \le k \le b_0, \\ k \equiv 1 \mod 2}} c_k\right) \wedge X_{m+1}^l. \end{split}$$
This shows the equivariance of $\tilde{\varphi}$.

This shows the equivariance of $\tilde{\varphi}$.

3.3. Injectivity. In the case of Theorem 1.2 we know that the dimension of $V^{n,d}$ agrees with the dimension of $\Lambda^{l}H_{1}(\Sigma_{g}^{1},\mathbb{C})/\omega \wedge \Lambda^{l-2}H_{1}(\Sigma_{g}^{1},\mathbb{C})$, so we can show that φ – which here denotes the map to the quotient - is an isomorphism by showing that it is surjective, which we will do next. This also shows the injectivity claimed in Theorem 1.1, by construction of $\tilde{\varphi}$.

Proof of Theorems 1.1 and 1.2. Denote by $\overline{\varphi}$ the composition of φ with the projection to the quotient $\Lambda^{l}H_{1}(\Sigma_{g}^{1},\mathbb{C})/\omega \wedge \Lambda^{l-2}H_{1}(\Sigma_{g}^{1},\mathbb{C})$. We just need to show that $\overline{\varphi}$ is injective, as we know that the spaces have the same dimension. Assume that v is in the kernel of $\overline{\varphi}$; then

$$e_i v = T_i v = -i(v + iT_i v) + iv,$$

so

$$\overline{\varphi}(e_i v) = -it_{c_i}\overline{\varphi}(v) + i\overline{\varphi}(v) = 0,$$

which shows that ker $\overline{\varphi}$ is a $\mathrm{TL}_n(\exp(-\pi i/4))$ -subrepresentation of $V^{n,d}$. But it follows from Corollary 4.8 of [RSA12] that the representation on $V^{n,d}$ is irreducible (as noted in the remarks on the case $\beta = 0$ following their Corollary), so ker $\overline{\varphi}$ is either 0 or $V^{n,d}$. It is easy to check that the latter is not the case, and so we obtain the Theorem. \square

Remark 3.5. We remark that the injectivity above can also be shown by explicitly constructing a basis $a_1, a_2, \ldots, a_i, b_1, b_2, \ldots, b_j$ of $\Lambda^l H_1(\Sigma, \mathbb{C})$ in such a way that φ surjects onto span $\{a_k\}$, $\omega \wedge \Lambda^{l-2}H_1(\Sigma, \mathbb{C}) = \operatorname{span}\{b_k\}$, and $\operatorname{span}\{a_k\}$ has the right dimension.

Proof of Theorem 1.3. This follows from Theorem 1.2 in the following way: we can define a map $h_1: V^{n,d} \to V^{n-1,d-1}$ that maps D to 0 if n is not connected to ∞ , and otherwise to the diagram given by removing the arc going from n to ∞ ; here we let $V^{n-1,-1} = V^{n-1,n+1} = \{0\}$. Likewise, we define a map $h_2: V^{n,d} \to V^{n-1,d+1}$ that maps a diagram to 0 if n is connected to ∞ , and otherwise to the diagram where the point connected to n is now connected to ∞ . For a diagram of the latter type, we define $g(D) = \alpha X_e$, where α is a scalar depending on D, and e is an arc such that $\varphi(D) = g(D) \wedge \varphi(h_2(D))$. It is clear that $g(D) = \alpha \sum_{i=k}^{n-1} c_i$ for some k, and by looking at the block decomposition of φ with respect to these two kinds of diagrams and with respect to the two subspaces $c_{n-1} \wedge \Lambda^{l-1} \operatorname{span}\{c_i \mid i = 1, 2, \ldots, n-2\}$ and its orthogonal complement (using wedge products of c_i 's as an orthogonal basis), we see that φ is injective, as it has the form

$$\left(\begin{array}{c|c} \varphi^{n-1,d-1} \circ h_1 & * \\ \hline 0 & \alpha c_{n-1} \wedge \varphi^{n-1,d+1} \circ h_2 \end{array}\right),$$

where the diagonal blocks are injective by Theorem 1.2; here, we interpret $\varphi^{n-1,-1}$ and $\varphi^{n-1,n+1}$ as the maps between two 0-dimensional spaces.

4. The AMU conjecture

We turn now to the proofs of Theorems 1.4 and 1.6. Recall that these are concerned with the genus 0 quantum SU(N)-representations $\rho_{N,k}^{\lambda}$, depending on a level k and a Young diagram $\lambda \in L_{N,k}$. We have taken as our definition of $\rho_{N,k}^{\lambda}$ that based on conformal field theory, so as to directly relate the quantum representations to those discussed in the previous sections, through the following result of Kanie; see [Kan89] and [Uen08]. We refer to [AU11, Sects. 4–7] for a complete description of the relevant skein theoretically defined quantum representations and the correspondence to the ones at hand.

Theorem 4.1 (Kanie). The Jones representation π_{λ} associated to a Young diagram $\lambda \in \Lambda_n^{(N,k+N)}$ is isomorphic at $q = \exp(\frac{2\pi i}{k+N})$ to the quantum SU(N)-representation $\rho_{N,k}^{T(\lambda)}$, where $T(\lambda)$ is defined in Section 1.2. In particular, the representation $\pi_q^{n,d}$ of B_n – with $q = \exp(\frac{2\pi i}{k+N})$ – is isomorphic to the quantum SU(N)-representation $\rho_{N,k}^{\lambda}$, where $\lambda = ((n+d)/2 \ge (n-d)/2)$ when N > 2, and $\lambda = (d)$ when N = 2.

Lemma 4.2. Let X_n denote the set of primitive n'th roots of unity. Then for every $z \in U(1)$, there exist $z_n \in X_n$ such that $\lim_{n\to\infty} z_n = z$.

Remark 4.3. For z = -1 – which in fact is the only case we will need in the proof of Theorem 1.4 – this is [AMU06, Lem. 5.1], in which the sequence z_n is constructed explicitly. We include this more general case for use in Appendix A below.

Proof of Lemma 4.2. Write $z = \exp(2\pi i\alpha), \alpha \in [0, 1)$, and let

$$Y_n = \{ m \mid 0 < m < n, \ \gcd(m, n) = 1 \}.$$

The largest gap between two consecutive points of Y_n is bounded from above by j(n), the ordinary Jacobsthal function at n. The result of Iwaniec [Iwa78] allows us to bound the size of this gap as $j(n) = O(\log^2(n))$. Thus, since $\log^2(n)/n \to 0$ as $n \to \infty$, we may choose $\alpha_n \in Y_n/n$ such that $\alpha_n - \alpha \to 0$ for $n \to \infty$. Letting $z_n = \exp(2\pi i \alpha_n) \in X_n$, we obtain the result since $z_n z^{-1} \to 1$ as $n \to \infty$.

For $\varphi \in M(0, n)_{\infty}$, define $\operatorname{sr}_d(\varphi) : \mathbb{R} \to \mathbb{R}_{>0}$ by

$$\operatorname{sr}_d(\varphi)(x) = \operatorname{sr}(\eta_{\exp(-\pi i x/4)}^{n,d}(\varphi)),$$

for $d \in \{0, ..., n\}$ with $d \equiv n \mod 2$, where sr denotes the spectral radius of a linear map.

Lemma 4.4. Let $\varphi \in M(0,n)_{\infty}$. If $\operatorname{sr}_d(\varphi)(x_0) > 1$ for some $x_0 \in [0,1]$, then there exists $k_0 \in \mathbb{N}$ such that the order of $\rho_{N,k}^{\lambda}(\varphi)$ is infinite for all $k > k_0$, where $\lambda = ((n+d)/2 \ge (n-d)/2)$.

Proof. It follows from Theorem 4.1 and the relations described in Section 2 that it suffices to show that for some k_0 , the order of $\eta_A^{n,d}(\varphi)$ is infinite for $q = A^4 = \exp(2\pi i/(k+N))$ for all k with $k > k_0$. On the other hand, the order of $\eta_A^{n,d}(\varphi)$ at a general primitive 4(k+N)'th root of unity $A = \exp(\frac{2\pi i l}{4(k+N)})$ is independent of l.

Now, apply Lemma 4.2 to $q = \exp(-\pi i x_0)$ to obtain primitive 4(k+N)'th roots of unity A_k with $A_k^4 \to q$. Since $\operatorname{sr}_d(\varphi)$ is continuous, $\operatorname{sr}(\eta_{A_k}^{n,d}(\varphi)) > 1$, for all sufficiently large k, and a linear map having an eigenvalue of absolute value greater than 1 necessarily has infinite order.

To show that a given pseudo-Anosov $\varphi \in M(0, n)_{\infty}$ has infinite order in $\rho_{N,k}^{\lambda}$ for all but finitely many levels, it thus suffices to find x_0 as above. As we will see below, by Theorems 1.2 and 1.3, this is possible for those pseudo-Anosovs whose stretch factors are given by their action on the homology of the double cover. In case d = n-2, the claims will follow immediately from Lemma 4.4 but in general, we will need the following remarks on the actions of surface diffeomorphisms on wedge products of homology.

Remark 4.5. Let f_* be the action on $H_1(\Sigma, \mathbb{C})$, where Σ is a surface of genus g with 0 or 1 boundary components, induced by a diffeomorphism f. Let $\lambda_1, \ldots, \lambda_{2g}$ be the diagonal entries in the Jordan normal form of the matrix for f_* . Then the action of f on $\Lambda^l H_1(\Sigma, \mathbb{C})$ will have an eigenvector of eigenvalue $\prod_{i \in I} \lambda_i$ for any subset $I \subseteq \{1 : 2g\}$ of size l, given by wedging together the corresponding vectors in the Jordan normal form, in such a way that a non-eigenvector is only included if all the preceding vectors in its block are also included. As all of these eigenvectors are of the form $\bigwedge_{i=1}^{l} v_i$, they can not be in the subspace $\omega \wedge \Lambda^{l-2}H_1(\Sigma, \mathbb{C})$ if $l \leq g$, and therefore they define eigenvectors of the same eigenvalue in the quotient $\Lambda^l H_1(\Sigma, \mathbb{C})/\omega \wedge \Lambda^{l-2} H_1(\Sigma, \mathbb{C})$.

Assume now that the action on $H_1(\Sigma, \mathbb{C})$ has an eigenvalue with absolute value strictly greater than 1. The action is symplectic, so the eigenvalues come in pairs λ, λ^{-1} , and there must be at least g columns in the Jordan normal form having diagonal entry with absolute value at least 1. Therefore the action on $\Lambda^l H_1(\Sigma, \mathbb{C})$ has an eigenvalue with absolute value greater than 1 for $l \leq g$, and by the above considerations, so does the action on $\Lambda^l H_1(\Sigma, \mathbb{C})/\omega \wedge \Lambda^{l-2} H_1(\Sigma, \mathbb{C})$.

Proposition 4.6. Let $\lambda \in L_{N,k}$, let $f \in M(0,n)_{\infty}$, and assume that the action \hat{f}_* of $\hat{f} = \Psi(f)$ on $H_1(\Sigma_g^m)$ has spectral radius strictly greater than 1. Then the order of $\rho_{N,k}^{\lambda}(f)$ is infinite for all but finitely many levels k.

Proof. As mentioned in the introduction, the main difficulty lies in showing the claim for the case where λ comes from a general 2-row diagram in $\Lambda_n^{(N,k+N)}$: as noticed above, the claim follows immediately from our main theorems in case d = n - 2, by application of Lemma 4.4 with $x_0 = 1$, since $\operatorname{sr}_d(f)(-1) = \operatorname{sr}(\hat{f}_*)$. This also implies the claim when λ comes from a hook by application of [FLW02, Thm. 2.2]; this theorem states that the Jones representations for hooks are equivalent (up to tensoring by particular characters) to exterior powers of the reduced Burau representation $\eta_A^{n,n-2}$, the power depending on the number of rows of the hook.

For general d, we also need to ensure that an appropriate eigenvector – of eigenvalue with absolute value strictly greater than 1 – is actually contained in the image of the morphism of representations. This, on the other hand, is guaranteed by Remark 4.5 in case n is odd.

Assume now that n is even and that d < n-2. Since \hat{f} preserves the boundaries pointwise, it defines a diffeomorphism of Σ_{g+1}^1 , obtained by gluing to Σ_g^2 a pair of pants. Denote by ι the induced map on (wedge products of) homology. On the level of diagrams, this corresponds to the inclusion $V^{n,d} \hookrightarrow V^{n+1,d+1}$ obtained by adding to a diagram the point n + 1 and connecting it to ∞ . This of course corresponds to the decomposition $V^{n+1,d+1} = V^{n,d} \oplus V^{n,d+2}$ as a vector space, as described in the proof of Theorem 1.3 (but note that n + 1 is now odd); that is, $V^{n,d+2}$ is spanned by diagrams where n + 1 is not connected to ∞ . Now, even though the action of f does not preserve the decomposition, it is clearly block triangular.

With these identifications, we have a diagram

which is commutative up to a power of -i. We wish to show that the action of f on $V^{n,d} \subseteq V^{n+1,d+1}$ contains eigenvectors of the appropriate eigenvalues. Suppose that the eigenvalues of the action of f_* on $H_1(\Sigma_a^2, \mathbb{C})$, counted with algebraic multiplicity, have absolute values

$$(x_1,\ldots,x_m,1,1,\ldots,1,x_m^{-1},\ldots,x_1^{-1})$$

with $x_i > 1$ for all i so that as before, $m \ge 1$. Consider first the case $l \le m$. As in Remark 4.5, the action of \hat{f}_* on $\Lambda^l H_1(\Sigma_g^2, \mathbb{C})$ has an eigenvector $v = v_1 \land \cdots \land v_l$ whose eigenvalue has absolute value $x = x_1 \cdots x_l$. Now, $\iota(v)$ is an eigenvector for the induced action with the same eigenvalue (up to the same root of unity), and it follows from the case of n odd that $\operatorname{Im}(\varphi^{n+1,d+1})$ contains an eigenvector which has the same eigenvalue as $\iota(v)$. Moreover, since the eigenvectors arising from $V^{n,d+2}$ all have absolute value strictly less than x (as we take only the (l-1)'st wedge products), we obtain the desired eigenvector of $\eta_{\exp(-\pi i/4)}^{n,d}(f)$. The conclusion now follows as in the case of odd n.

The case l > m is similar but involves also a small combinatorial exercise as in this case, there may also be eigenvectors coming from $V^{n,d+2}$ with eigenvalue of absolute value x. Let $d_x^{n,d}$ be the

sum of the algebraic multiplicities of eigenvalues of absolute value x of the action of f on $V^{n,d}$. We claim that for n even,

(3)
$$d_x^{n,d} = \binom{2g+1-2m}{l-m} - \binom{2g+1-2m}{l-m-2}.$$

To see this, we appeal again to the decomposition used above, as a simple extension of the argument from Remark 4.5 shows that

$$d_x^{n+1,d+1} = \binom{2g+2-2m}{l-m} - \binom{2g+2-2m}{l-m-2}.$$

Now, since $d_x^{n+1,d+1} = d_x^{n,d} + d_x^{n,d+2}$, equation (3) follows by induction on l, starting at l = m, by using well-known recursive formulas for binomial coefficients. Since $d_x^{n,d} > 0$, this completes the proof.

Proof of Theorem 1.4. Let $f \in M(0,n)_{\infty}$ be a homological pseudo-Anosov, let \hat{f} denote its image in M(0, n + 1), and let $\hat{f} \in \overline{M}(g, m)$ denote the image of f under the Birman-Hilden map used in Theorems 1.2 and 1.3 with the appropriate values of g and m. Everything has been set up so that \hat{f} is a pseudo-Anosov of Σ_g^m with the same stretch factor as \tilde{f} , and that, moreover, \hat{f} has *orientable* invariant foliations. This follows by the exact same reasoning as in the similar setup in Theorem 5.1 of [BB07]. In short: the orientability of a foliation is determined by the vanishing of its associated orientation homomorphism, and this on the other hand is ensured by the assumptions on the degrees of the singularities.

Now, the stretch factor of any pseudo-Anosov with orientable invariant foliations is simply the spectral radius of its action on homology which is therefore strictly greater than 1. This is a well-known result and in fact a criterion for having orientable foliations; see e.g. [BB07, Lemma 4.3] and the discussion preceding it.

The claim then follows directly from Proposition 4.6.

Proof of Theorem 1.6. Consider now the case where N = 2, n is even, and λ is the empty diagram. Here, the level k quantum SU(2)-representation, rescaled on each generator by a suitable k-dependent root of unity, defines a representation of $\widetilde{M}(0, n)$, equivalent by construction to $\rho_{2,k}$ (see [Jon87, Sect. 10]). As multiplication by a root of unity does not change whether or not the order of a linear map is finite or infinite, we obtain from Theorem 1.4 the claimed result.

Remark 4.7. More generally, in [Jon87, Sect. 10], Jones finds that his representations may be tweaked by roots of unity to descend to the mapping class groups of spheres whenever the associated Young diagram λ is rectangular. Thus, the same is of course true for the quantum SU(N)representations $\rho_{N,k}^{\lambda}$. One could therefore proceed as in [AMU06, Sect. 4], define new quantum representations for mapping class groups of punctured spheres, and immediately obtain a version of Theorem 1.6 for those.

Example 4.8. At this point it may be worth remarking that examples of homological pseudo-Anosovs are plenty.

- In [LT11, App. A], the authors provide several examples of homological (as well as non-homological) braids, some of which we shall return to in the appendix below.
- On a closed torus, the stretch factor of a pseudo-Anosov is always given by its action on homology, and so we recover the main result of [AMU06].
- As noted in the proof of Theorem 1.4, [BB07, Lemma 4.3] tells us that if the spectral radius of the action of a pseudo-Anosov on homology equals its stretch factor, the pseudo-Anosov must necessarily have orientable invariant foliations, and so we can appeal directly to any of the existing homological constructions of pseudo-Anosovs to obtain interesting examples. One such family of examples arises as a special case of the pseudo-Anosovs described in [Pen88] (see also [BC88, Sect. 5]) on the level of the covering surfaces, which passing through the Birman–Hilden homomorphism may be described as follows. Suppose that n is even. Then any word in the generators $\sigma_1, \ldots, \sigma_{n-1}$ such that the sign of the exponents

for odd-indexed generators all agree, such that exponents of even-indexed generators all have the opposite sign as the odd-indexed ones, and such that each generator appears at least once, is a homological pseudo-Anosov.

4.1. Determining stretch factors. In [AMU06, Cor. 5.8], the authors go on to show that the stretch factor of any given pseudo-Anosov of a four times punctured sphere may be obtained as limits of eigenvalues of the quantum representations of φ . The analogous statement in our general case is the following.

Corollary 4.9. For any homological pseudo-Anosov $\varphi \in M(0,n)_{\infty}$ and Young diagram λ as in Section 1.2, there exist eigenvalues λ_k , $\tilde{\lambda}_k$ of $\rho_{N,k}^{\lambda}(\varphi)$ such that $\sqrt{|\lambda_k \tilde{\lambda}_k|}$ tends to the stretch factor of φ as $k \to \infty$.

Proof. The statement follows from the proof of Proposition 4.6 by continuity in A of the eigenvalues of $\eta_A^{n,d}(\varphi)$, as these include at $q = A^4 = -1$ the values $\tau \mu$ and $\tau \mu^{-1}$, where $|\tau|$ is the stretch factor of φ (and likewise, μ is a product of eigenvalues of the induced action on homology).

APPENDIX A. EXAMPLES, EXPERIMENTS, AND OBSERVATIONS

A.1. A pseudo-Anosov in the Torelli group. The preceding discussion begs the question of the behaviour of those pseudo-Anosov mapping classes for which the action on homology contains no information. In particular, we could consider examples of pseudo-Anosovs acting trivially on the homology of the double cover. That pseudo-Anosovs in the Torelli group, i.e. the kernel of the action of Γ_g on $H_1(\Sigma_g)$, exist for $g \ge 2$ was shown in [Thu88]. One concrete such element was constructed by Brown [Bro03, App.] for g = 2, and appealing once again to the setup of the Birman-Hilden theorem, this allows us to obtain the desired braid $\varphi \in B_6$.

More precisely, define

(4)
$$\varphi = \eta \xi^{-1}$$

explicitly in terms of the standard braid group generators by

$$\begin{split} \xi &= \gamma \eta \gamma^{-1}, \\ \eta &= \sigma_4^2 \sigma_5 \sigma_4^2 \sigma_5^2 \sigma_4^2 \sigma_5 \sigma_4^2, \\ \delta &= \sigma_4 \sigma_5 \sigma_3 \sigma_2 \sigma_1 \sigma_2^{-1} \sigma_3^{-1} \sigma_5^{-1} \sigma_4^{-1}, \\ \gamma &= \delta \sigma_3 \sigma_2^{-1} \sigma_1^{-1} \delta \sigma_1 \sigma_2 \sigma_3^{-1} \delta^{-1}. \end{split}$$

As we have seen, to show that a pseudo-Anosov $\psi \in B_n$ has infinite order in the quantum representations for all but finitely many levels, it suffices to show that the function $\operatorname{sr}_d(\psi) : [0,1] \to \mathbb{R}_{>0}$ given by

$$\operatorname{sr}_d(\psi)(x) = \operatorname{sr}(\eta_{\exp(-\pi i x/4)}^{n,d}(\psi))$$

is greater than 1 for some value $x \in [0, 1]$. The result of evaluating this function for the pseudo-Anosov φ of (4) is shown¹ in Figure 8. We see for instance that $\operatorname{sr}_0(\varphi)(x) = 1$ for $x \in [0, \frac{1}{2}] \cup \{1\}$ but that this is not the case in general.

Proposition A.1 (Computer assisted). There exist pseudo-Anosovs in ker $\eta_{\exp(-\pi i/4)}^{n,d}$ that have infinite order in the corresponding $\rho_{N,k}^{\lambda}$ for all but finitely many k.

Previously, we saw that sr naturally determines the stretch factors of homological pseudo-Anosov but it is not clear in which sense, if any, this is true for the mapping class φ above.

¹This figure, and all other figures in this section, were created using Sage 5.13 and Mathematica 9.0.



FIGURE 8. Graphs of $\operatorname{sr}_d(\varphi)(x)$ for $x \in [0, 1]$.

A.2. Infinite order at low levels. Other examples exhibiting similar behaviours include the stretch factor minimizing pseudo-Anosov braids of [LT11]; plots of sr are shown for these in Figure 9. In each of these, we let $\psi_n = \sigma_1 \cdots \sigma_n$, and the dashed line indicates the stretch factor of the pseudo-Anosov in question. In particular, the braid in Figure 9b provides an interesting example: it has finite k-dependent order in $\rho_{N,k}^{\lambda}$ for every level k with N > 2 and $\lambda = \square$. We remark that in [AMU06, Conjec. 2.4], the authors do not mention any possible assumptions on the labelling in the relevant quantum representations.

Notice also that these graphs allow us to deduce more precisely at *which* levels k, the orders of the mapping classes in question are infinite. For example, in Figure 10, we consider again the braid of Figure 9b, now viewed as an element of B_6 and highlight the possible specializations of A for the level k quantum SU(2)-representation when $k = 5, \ldots, 12$.

From this figure alone, we obtain the following.

Proposition A.2. The level 8 quantum SU(2)-representation, with λ the empty Young diagram, of the mapping class $\sigma_1 \sigma_2 \sigma_3^{-1} \in B_6$ is infinite.

Proof. Whereas the claim follows from Figure 10, let us also note that one finds by explicit calculation – requiring no electronic aid – that for a particular ordering in our preferred basis,

$$\eta_A^{6,0}(\sigma_1\sigma_2\sigma_3^{-1}) = A \begin{pmatrix} 0 & A^{-6} & 0 & -1 & A^2 \\ A^2 & -1 + A^{-4} & -1 & 0 & 0 \\ 0 & 0 & 0 & A^{-6} - A^{-2} & 1 \\ 0 & 0 & -A^{-2} & -1 + A^{-4} & A^2 \\ 0 & 0 & 0 & -A^2 & A^4 \end{pmatrix}$$

Substituting the primitive $4 \cdot (8+2)$ 'th root of unity $A = \exp(2\pi i \frac{3}{40})$, one obtains that

$$\operatorname{sr}(\eta_A^{6,0}(\sigma_1\sigma_2\sigma_3^{-1})) \approx 1,665 > 1.$$

While we have not considered higher genus quantum representations of mapping class groups in the present paper, let us notice that it follows from Proposition A.2 and a standard factorization argument that the quantum representations of closed surfaces of genus g > 2 have infinite image at level k = 8; this extends an earlier result due to Masbaum [Mas99] (which is indeed the reason



FIGURE 9. Plots of spectral radii for braids with small stretch factors.

we considered this particular level). In fact, at level k = 8, the image of the quantum SU(2)representation obtained from the sphere with four marked points is known to be finite by the result of [LPS13], and so one can not use the Jones representations to obtain a similar result for g = 2, k = 8. However, the second author has found – again aided by computer calculation – example mapping classes whose quantum representations have infinite order for g = 2, k = 8, by embedding in the genus 2 surface a one-holed torus rather than a four-holed sphere; see e.g. [Jør13, Prop. 4.16], [Jør11, p. 49].

Now, the question about the size of the image is still open for levels k = 1, 2, and 4, and casting a sidelong glance at Figures 9e and 9f, noting that at the fourth root of unity $q = A^4 = i$, the function sr is greater than 1, one could be inclined to believe that these provide infinite order examples for k = 2. Recall however that these specializations do not correspond to the level 2 quantum representations in which one finds that the two elements have orders 28 and 16



FIGURE 10. The graph of sr₀ for $\sigma_1 \sigma_2 \sigma_3^{-1} \in B_6$.

respectively. Indeed, in the cases k = 1, 2, and 4, the only possible specializations of values of q allowed by skein theory are $q = \exp(2\pi i/(k+2))$. The corresponding representations are known to be unitary and so no information may be gained from considering spectral radii.

A.3. Bigelow's element of the kernel of the Burau representation. As a final remark, the connection between the various sr_d is an interesting question. As a single example, Figure 11 shows $\operatorname{sr}_d(\psi)$ for Bigelow's [Big99] element $\psi \in \operatorname{ker}(\pi_q^{5,3})$, given explicitly by

 $\psi = [\psi_1^{-1} \sigma_4 \psi_1, \psi_2^{-1} \sigma_4 \sigma_3 \sigma_2 \sigma_1^2 \sigma_2 \sigma_3 \sigma_4 \psi_2],$

where

$$\begin{split} \psi_1 &= \sigma_3^{-1} \sigma_2 \sigma_1^2 \sigma_2 \sigma_4^3 \sigma_3 \sigma_2, \\ \psi_2 &= \sigma_4^{-1} \sigma_3 \sigma_2 \sigma_1^{-2} \sigma_2 \sigma_1^2 \sigma_2^2 \sigma_1 \sigma_4^5. \end{split}$$

References

- [And06] Jørgen Ellegaard Andersen. Asymptotic faithfulness of the SU(n) representations. Annals of Mathematics, 163:347–368, 2006.
- [AU07a] Jørgen Ellegaard Andersen and Kenji Ueno. Abelian Conformal Field theories and Determinant Bundles. International Journal of Mathematics, 18:919–993, 2007.
- [AU07b] Jørgen Ellegaard Andersen and Kenji Ueno. Constructing modular functors from conformal field theories. Journal of Knot theory and its Ramifications, 16(2):127–202, 2007.
- [AU11] Jørgen Ellegaard Andersen and Kenji Ueno. Construction of the Reshetikhin-Turaev TQFT from conformal field theory. http://arxiv.org/pdf/1110.5027, 2011.
- [AU12] Jørgen Ellegaard Andersen and Kenji Ueno. Modular functors are determined by their genus zero data. Quantum Topology, 3:255–291, 2012.
- [BB07] Gavin Band and Philip Boyland. The Burau estimate for the entropy of a braid. Algebr. Geom. Topol., 7:1345–1378, 2007. ISSN 1472-2747; 1472-2739/e.
- [BC88] Steven A. Bleiler and Andrew J. Casson. Automorphisms of Surfaces after Nielsen and Thurston. Cambridge University Press, 1988.



FIGURE 11. The functions $\operatorname{sr}_d(\psi)(x), x \in [0,1]$, for a $\psi \in \operatorname{ker}(\pi_q^{5,3})$.

[BH73] Joan S. Birman and Hugh M. Hilden. On Isotopies of Homeomorphisms of Riemann Surfaces. The Annals of Mathematics, Second Series, 97(3):424–439, 1973.

- [BHMV92] Christian Blanchet, Nathan Habegger, Gregor Masbaum, and Pierre Vogel. Three-manifold invariants derived from the Kauffman bracket. *Topology*, 31:685–699, 1992.
- [BHMV95] Christian Blanchet, Nathan Habegger, Gregor Masbaum, and Pierre Vogel. Topological quantum field theories derived from the Kauffman bracket. *Topology*, 34:883–927, 1995.
- [Big99]Stephen J. Bigelow. The Burau representation is not faithful for n = 5. Geom. Topol., 3:397-404, 1999.[Bla00]Christian Blanchet. Hecke algebras, modular categories and 3-manifolds quantum invariants. Topology, 39(1):193-223, 2000.
- [Bro03] Richard J. Brown. Generating quadratic pseudo-Anosov homeomorphisms of closed surfaces. Geom. Dedicata, 97:129–150, 2003.
- [FLW02] Michael H. Freedman, Michael J. Larsen, and Zhenghan Wang. The two-eigenvalue problem and density of Jones representation of braid groups. *Commun. Math. Phys.*, 228(1):177–199, 2002. ISSN 0010-3616; 1432-0916/e.
- [FM11] Benson Farb and Dan Margalit. A primer on mapping class groups. Princeton Mathematical Series. Princeton, NJ: Princeton University Press. xiv, 492 p., 2011.
- [FWW02] Michael Freedman, Kevin Walker, and Zhenghan Wang. Quantum SU(2) faithfully detects mapping class groups modulo center. Geom. Topol., 6:523–539, 2002.

[Hit90] Nigel Hitchin. Flat Connections and Gometric Quantization. Commun. Math. Phys., 131:347–380, 1990.

[Iwa78] Henryk Iwaniec. On the problem of Jacobsthal. Demonstr. Math., 11:225–231, 1978. ISSN 0420-1213.
 [Jon83] Vaughan F.R. Jones. Braid groups, Hecke algebras and type II₁ factors. Geometric methods in operator

algebras, Proc. US-Jap. Semin., Kyoto/Jap. 1983., 1983.

- [Jon87] Vaughan F.R. Jones. Hecke algebra representations of braid groups and link polynomials. Ann. Math., 126(2):335–388, 1987.
- [Jør11] Søren Fuglede Jørgensen. Quantum representations of mapping class groups, progress report, 2011. Available at http://maths.fuglede.dk.
- [Jør13] Søren Fuglede Jørgensen. Semiclassical properties of the quantum representations of mapping class groups. PhD thesis, Aarhus University, 2013. Available at http://maths.fuglede.dk.
- [Kan89] Yukihiro Kanie. Conformal Field Theory and the Braid Group. Bulletin of the Faculty of Education Mie University, 40:1–43, 1989.
- [Kas01] Yasushi Kasahara. An expansion of the Jones representation of genus 2 and the Torelli group. Algebr. Geom. Topol., 1:39–55, 2001.
- [Kob12] Thomas Koberda. Asymptotic linearity of the mapping class group and a homological version of the Nielsen-Thurston classification. Geom. Dedicata, 156:13–30, 2012. ISSN 0046-5755; 1572-9168/e.
- [LPS13] Yves Laszlo, Christian Pauly, and Christoph Sorger. On the monodromy of the Hitchin connection. J. Geom. Phys., 64:64–78, 2013.
- [LT11] Erwan Lanneau and Jean-Luc Thiffeault. On the minimum dilatation of braids on punctured discs. Geom. Dedicata, 152:165–182, 2011.

- [Mas99] Gregor Masbaum. An element of infinite order in TQFT-representations of mapping class groups. Contemp. Math, pages 137–139, 1999.
- [Mas08] Gregor Masbaum. On representations of mapping class groups in Integral TQFT. Oberwolfach Reports, 5:1157–1232, 2008.
- [McM] Curtis McMullen. Entropy on Riemann surfaces and Jacobians of finite covers. Comment. Math. Helv., to appear.
- [Pen88] Robert Penner. A Construction of Pseudo-Anosov Homeomorphisms. Transactions of the American Mathematical Society, 310:179–197, 1988.
- [RSA12] David Ridout and Yvan Saint-Aubin. Standard Modules, Induction and the Structure of the Temperley-Lieb algebra, 2012. http://arxiv.org/abs/1204.4505v2.
- [RT90] Nicolai Yu. Reshetikhin and Vladimir G. Turaev. Ribbon graphs and their invariants derived from quantum groups. Comm. Math. Phys, 127:1–26, 1990.
- [RT91] Nicolai Yu. Reshetikhin and Vladimir G. Turaev. Invariants of 3-manifolds via link polynomials and quantum groups. Invent. Math., 103(3):547-597, 1991. CODEN INVMBH. ISSN 0020-9910. URL http: //dx.doi.org/10.1007/BF01239527.
- [San12] Ramanujan Santharoubane. Limits of the quantum SO(3) representations for the one-holed torus. J. Knot Theory Ramifications, 21(11):13, 2012. ISSN 0218-2165.
- [Thu88] William P. Thurston. On the geometry and dynamics of diffeomorphisms of surfaces. Bull. Amer. Math. Soc. (N.S.), 19(2):417–431, 1988.
- [Tur10] Vladimir G. Turaev. Quantum invariants of knots and 3-manifolds, volume 18 of de Gruyter Studies in Mathematics. Walter de Gruyter & Co., Berlin, revised edition, 2010. ISBN 978-3-11-022183-1. xii+592 pp. URL http://dx.doi.org/10.1515/9783110221848.
- [TUY89] Akihiro Tsuchiya, Kenji Ueno, and Yasuhiko Yamada. Conformal field theory on universal family of stable curves with gauge symmetries. In *Integrable systems in quantum field theory and statistical mechanics*, volume 19 of Adv. Stud. Pure Math., pages 459–566. Academic Press, Boston, MA, 1989.
- [Uen08] Kenji Ueno. Conformal field theory with gauge symmetry, volume 24 of Fields Institute Monographs. American Mathematical Society, Providence, RI, 2008. ISBN 978-0-8218-4088-7. viii+168 pp.
- [Wan10] Zhenghan Wang. Tolological quantum computation. Providence, RI: American Mathematical Society (AMS), 2010. ISBN 978-0-8218-4930-9/pbk. xiii + 115 pp.
- [Wen88] Hans Wenzl. Hecke algebras of type A_n and subfactors. *Invent. Math.*, 92(2):349–383, 1988. ISSN 0020-9910; 1432-1297/e.
- [Wit89] Edward Witten. Quantum field theory and the Jones polynomial. Comm. Math. Phys., 121:351–399, 1989.

Centre for Quantum Geometry of Moduli Spaces, Faculty of Science, Aarhus University, DK-8000 Aarhus C, Denmark

E-mail address: jk@qgm.au.dk

DEPARTMENT OF MATHEMATICS, BOX 480, UPPSALA UNIVERSITY, SE-75106 UPPSALA, SWEDEN *E-mail address*: soren.fuglede.jorgensen@math.uu.se

20